

Maths AA HL Questions:

Question 1:

[Maximum mark: 4]

$$\text{Solve } \tan(2x-5^\circ) = 1$$

[for $0 < x < 180^\circ$]

Answer Q1:

Since $\tan(x)$ is only 1 when the terminal angle is 45° , the angles used for $(2x-5)$ are 45° and 225° .

Hence, the answers are 25° **and** 115° .

Question 2:

[Maximum mark: 5]

$$\text{Solve } 3 \cdot 9^x + 5 \cdot 3^x - 2 = 0$$

Answer Q2:

Convert 9^x to 3^{2x} and then substitute 3^x into the equation. You will then get $3u^2 + 5u - 2 = 0$.

When solving the quadratic, you will get $u = 1/3$ and $u = -2$.

Then you substitute $u = 3^x$ back into the equation.


Since you cannot get -2 from an exponential function, you can eliminate -2.

Therefore, since $3^{-1} = 1/3$, the only answer is $x = -1$.

Question 3:

[Maximum mark:

The following diagram shows quadrilateral OABC.



Quadrilateral OABC is symmetrical across line OB.

Point C is $(3, 3\sqrt{3})$ and Point A is $(6, 0)$.

- Find the midpoint of A and C.
- The equation of the line OB.

Given that AB is perpendicular to OA,

- Find the area of quadrilateral OABC.

Answer Key:

- The average of 3 and 6 is 4.5. The average of 0 and $3\sqrt{3}$ is $(3\sqrt{3})/2$. The point is $(4.5, (3\sqrt{3})/2)$.
- The gradient of line AC is $-\sqrt{3}$. AC and OB are perpendicular since OB is a line of symmetry. Hence, the gradient of line OB is $1/\sqrt{3}$. The intercept is 0, so no need to add anything.
- OB is a line of symmetry, so find the area of triangle OAB and multiply by 2. Since the gradient of line OB is $1/\sqrt{3}$, the point B is $(6, 2\sqrt{3})$. Therefore, the distance AB is $2\sqrt{3}$. By using the formula $0.5 * (b * h)$, the area OABC is $12\sqrt{3}$.

Question 4:

A species of bird has a different chance of nesting in different species.

In spring, the bird has a k chance of nesting.

In summer, the bird has a $k/2$ chance of nesting.

[Maximum mark: 6]
A bag contains 7 blue and 5 red marbles. Two marbles are selected at random without replacement.

1. Complete the tree diagram below. [3]

1. Find the probability that exactly one of the selected marbles is blue. [3]

a) Complete the tree diagram.

The probability of a bird not nesting in spring and not nesting in summer is $5/9$.

b) Show that $9k^2 - 27k + 8 = 0$.

c) $9k^2 - 27k + 8$ is fulfilled by $k = 1/3$ and $k = 8/3$. Explain why $k = 8/3$ is not valid.

ANSWER:

a) The leftmost box should have $1-k$. Both boxes on the right should have $1 - k/2$.

b) $(1 - k)(1 - k/2) = 5/9$ then $(1 - 3k/2 + k^2/2) = 5/9$ then $(2 - 3k + k^2) = 10/9$
then $18 - 27k + 9k^2 = 10$ then $9k^2 - 27k + 8 = 0$

c) If $k = 8/3$ then the probability exceeds 1 which is not possible.

Question 5:

$$f(x) = \frac{2(x+3)}{3(x+2)}$$

- a) What is the equation of the horizontal asymptote of the function?

$$g(x) = mx + 1, m \neq 0$$

- b) If $m > 0$, how many solutions to $f(x) = g(x)$ are there?
c) For what value of m will there only be one solution?
d) What is the range of values for m for which $f(x) = g(x)$ has two solutions **when** $x \geq 0$?

ANSWER:

- a) $f(x) = \frac{2x+6}{3x+6}$, so the horizontal asymptote is at $y = \frac{2}{3}$.
b) There will be two solutions.
c) $-1/6$
d) ?

ii) asks for what value of m will there be only one solution

I got it: I think its that the line $mx + 1$ is tangent to the curve so derivate. prove?

Question 6:

A farmer grows two different kinds of apples.

	Mean (g)	Standard Deviation
eating	100	20
cooking	140	40

For this question, you may assume that there are 95% of apples within two standard deviations.

- a) What is the percentage of eating apples that weigh more than 140g?

This farmer grows 80% eating apples.

In addition, a sorter sorts all apples that weigh more than 140g into one container.

- b) If you were to pick an apple out of this container, what is the probability that the apple you picked is an eating apple? Give your answer in $\frac{c}{d}$ where c and d are integers.

ANSWER:

- a) 5% of apples are outside two STDEVs. Half of these apples are on the larger side. 2.5%

- b) Cooking apples $> \frac{1}{2} \times \frac{1}{5} = \frac{1}{10} = \frac{10}{100}$, Eating apples $> \frac{1}{40} \times \frac{4}{5} = \frac{2}{100}$, so out of 100 apples, 12 of them will be greater than 140 grams and 2 of those 12 are eating apples.

Therefore, in simplest terms, the probability is $\frac{1}{6}$ where c is 1 and d is 6.

Question 7:

Equation 1 is $2x^3 - 7x^2 + \dots$

- a) What is the sum of the 3 roots [ABC] of this equation?

Equation 2 is $2z^5 - 11z^4 + \dots - 20$. Roots [ABC] in Equation 1 are also in Equation 2.

$h(z) = 0$ and $z = p + 3i$.

- b) Show that $p = 1$.

It is given that $h(0.5) = 0$. $A < B$.

- c) Find the value of AB .
d) Find the value of A and B separately.

ANSWER:

- a) $\frac{-b}{a}$ gives you $\frac{7}{2}$.

- b) $\frac{-b}{a}$ gives you $\frac{11}{2}$. Since there are 5 roots in Equation 2 and ABC are three of the roots, the last two roots have a sum of 2. In addition, since $z = p + 3i$ and all the coefficients are real, you can use CRT to say that $(p - 3i)$ and $(p + 3i)$ are the last two roots. Hence, you can say that $2p = 2$ and therefore $p = 1$.

- c) The product of all five roots is $\frac{20}{2} = 10$. Since $h(0.5) = 0$, you can set root C as 0.5.

Multiplying $(1 + 3i)$ and $(1 - 3i)$, you get 10. So, $AB * 0.5 * 10 = 10$. Therefore, $AB = 2$.

- d) Since both A and B are integers and B is greater than A , $A = 1$ and $B = 2$.

Question 8:

$$\text{Find } \lim_{h \rightarrow 0} \frac{\sec^4(x) - \cos^2(x)}{x^4 - x^2}.$$

ANSWER:

-3?

Question 9:

A teacher, for safety, has to split n students into two groups.

The first group must have **exactly** 3 people. The second group must have **at least** 3 people.

- a) Find the expression for the number of ways the groups can be sorted.

Two students can't work together and must be in different groups.

- b) The number of ways the groups can be sorted is halved. Find the number of students n .

ANSWER:

?

-----Section B-----

Question 10:

[Maximum mark: 16]

An arithmetic sequence begins a, p, q .

a) Show that $2p - q = a$.

A geometric sequence begins a, s, t .

b) Show that $s^2 = at$.

It is given that $q = t = 1$ and $a \neq 0$.

c) Show that $p > \frac{1}{2}$.

It is given that $a = 9, s > 0$ and $q = t = 1$.

d) List the first four terms of the arithmetic sequence.

e) List the first four terms of the geometric sequence.

Another arithmetic sequence begins $9 + \ln(9), 5 + \ln(3), \dots$

f) Find the common difference in this sequence.

g) Show that the sum of the first 10 terms is $-90 - 25 \ln(3)$.

ANSWER:

a) Let $p = a + d$ and $q = a + 2d$. Then substitute, $(2a + 2d) - (a + 2d) = a$.

b) Let $s = ar$ and $t = ar^2$. Then substitute, $a^2 r^2 = a \times ar^2$.

c) $2p - 1 = a$, then $p = \frac{a+1}{2}$. Since a cannot be 0 or negative, $p > \frac{1}{2}$.

d) With simple arithmetic, $d = -4$. Therefore, it goes 9, 5, 1, -3.

e) With simple arithmetic, $r = \frac{1}{3}$. Therefore, it goes 9, 3, 1, $\frac{1}{3}$.

f) $(5 + \ln(3)) - (9 + \ln(9)) = (5 + \ln(3)) - (9 + 2\ln(3)) = -4 - \ln(3)$.

g) Use the formula booklet. You get the right answer.

Question 11:

[Maximum mark: 19]

$\pi_1: 2x + 6y - 2z = 5$ and point A is $(2, \frac{1}{2}, 1)$

- a) Show that point A lies in π_1 .

$\pi_2: (k^2 - 6)x + (2k + 3)y + pz = q$, and $p = -6$.

- b) Given that both planes are perpendicular and A lies in Plane 2, find k and q .

From part c onwards, Plane 1 and Plane 2 are parallel. It is also given that $k=3$.

- c) Find p .

It is given that $q = \frac{51}{2}$.

- d) A line perpendicular to Plane 1 that goes through A meets Plane 2 at a point B. Find B.

- e) ?

ANSWER:

Answer currently incorrect, as point A was changed.

- a) Substitute point A in the equation for Plane 1. It should come out equal.

- b) Dot $(2, 6, -2)$ with $((k^2 - 6), (2k + 3), -6)$ in order to get a long equation. This equation should equal zero. Simplifying gives $k^2 + 6k + 9 = 0$ which is just $(k + 3)^2$. Therefore, $k = -3$. Substitute the k value and the coordinates for A into the equation and get $q = \frac{3}{2}$.

- c) Since Plane 1 and Plane 2 are parallel, the cartesian equation should be scalar multiples of each other. Therefore, since $k = 3$ and hence $x = 3$ and $y = 9$, p should be -3 .

- d) The resultant vector line is $(2, \frac{1}{2}, 1) + \lambda(2, 6, -2)$. Use the parametric form of this equation and substitute the variables into Plane 2's equation. Rearranging the equation gives

- e)

- f) Rearranging the equation gives $\lambda = \frac{25}{44}$. Use this value with the parametric equations to find

point B $(\frac{138}{44}, \frac{172}{44}, \frac{-94}{44})$.

- g) ?

Question 12:

maclaurin series with convergence and an induction

Original function $f(x) = (1-ax)^{-1/2}$

A: induction of the derivatives of a function

B: find the maclaurin of some function in A

C: show that it converges to..?

D: using $x = 0.1$, show the of sqrt3

Hence, show

$$(-4x + 1)^{-\frac{1}{2}} (-2x + 1)^{-\frac{1}{2}} \approx \frac{19x^2 + 6x + 2}{2}$$

Answer Q12:

(using proof by induction of the derivatives of a function)

SL:

Question 1: looking at the graph the questions were:

- a) i) $f(4)$
 - ii) $f(f(4))$
 - iii) $f^{-1}(3)$
- b) sketch $f^{-1}(x)$

Answer Q1:

Question 2: [Maximum mark: 4]

$$\text{Solve } \tan(2x-5^\circ) = 1$$
$$[\text{for } 0 < x < 180^\circ]$$

Answer Q2:

Since $\tan(x)$ is only 1 when the terminal angle is 45° , the angles used for $(2x-5)$ are 45° and 225° .
Hence, the answers are 25° and 115° .

Question 3:

Answer Q3:

Question 4:[Maximum mark: 5]

$$\text{Solve } 3 \cdot 9^x + 5 \cdot 3^x - 2 = 0$$

Answer Q4:

a) $9m^2+5m=0$ solve for m

b) Convert 9^x to 3^{2x} and then substitute 3^x into the equation. You will then get $3u^2 + 5u - 2 = 0$.

When solving the quadratic, you will get $u = 1/3$ and $u = -2$.

Then you substitute $u = 3^x$ back into the equation.

Since you cannot get -2 from an exponential function, you can eliminate -2 .

Therefore, since $3^{-1} = 1/3$, the only answer is $x = -1$.

Question 5: A species of bird has a different chance of nesting in different species.


In spring, the bird has a k chance of nesting.

In summer, the bird has a $k/2$ chance of nesting.

[Maximum mark: 6]

A bag contains 7 blue and 5 red marbles. Two marbles are selected at random without replacement.

1. Complete the tree diagram below. [3]



1. Find the probability that exactly one of the selected marbles is blue. [3]

- d) Complete the tree diagram.

The probability of a bird not nesting in spring and not nesting in summer is $\frac{5}{9}$.

- e) Show that $9k^2 - 27k + 8 = 0$.
- f) $9k^2 - 27k + 8$ is fulfilled by $k = \frac{1}{3}$ and $k = \frac{8}{3}$. Explain why $k = \frac{8}{3}$ is not valid.

Answer Q5:

ANSWER:

- d) The leftmost box should have $1-k$. Both boxes on the right should have $1 - \frac{k}{2}$.
- e) $(1 - k)(1 - \frac{k}{2}) = \frac{5}{9}$ then $(1 - \frac{3k}{2} + \frac{k^2}{2}) = \frac{5}{9}$ then $(2 - 3k + k^2) = \frac{10}{9}$
then $18 - 27k + 9k^2 = 10$ then $9k^2 - 27k + 8 = 0$
- f) If $k = \frac{8}{3}$ then the probability exceeds 1 which is not possible.

Question 6: $f(x) = \frac{2(x+3)}{3(x+2)}$

- e) What is the equation of the horizontal asymptote of the function?

$g(x) = mx + 1, m \neq 0$

- f) If $m > 0$, how many solutions to $f(x) = g(x)$ are there?
- g) For what value of m will there only be one solution?
- h) What is the range of values for m for which $f(x) = g(x)$ has two solutions **when** $x \geq 0$?

Answer Q6: ANSWER:

e) $f(x) = \frac{2x+6}{3x+6}$, so the horizontal asymptote is at $y = \frac{2}{3}$.

f) There will be two solutions.

g) $-1/6$

h) ?

ii) asks for what value of m will there be only one solution

I got it: I think its that the line $mx + 1$ is tangent to the curve so derivate. prove?

Question 7:

Answer Q7:

Question 8:

Answer Q8:

Question 9:

Answer Q9: